

ARMY RESEARCH LABORATORY



Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields

P. J. Stephens

F. J. Devlin

UNIVERSITY OF SOUTHERN CALIFORNIA

C. F. Chabalowski

U.S. ARMY RESEARCH LABORATORY

M. J. Frisch

LORENTZIAN INC.



ARL-TR-695

February 1995

19950222 025

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.

NOTICES

Destroy this report when it is no longer needed. DO NOT return it to the originator.

Additional copies of this report may be obtained from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The use of trade names or manufacturers' names in this report does not constitute endorsement of any commercial product.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE February 1995		3. REPORT TYPE AND DATES COVERED Final, January 1993-August 1994
4. TITLE AND SUBTITLE Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields			5. FUNDING NUMBERS PR: 1L161102AH43	
6. AUTHOR(S) P. J. Stephens,* F. J. Devlin,* C. F. Chabalowski, and M. J. Frisch**				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory ATTN: AMSRL-WT-PC Aberdeen Proving Ground, MD 21005-5066			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory ATTN: AMSRL-OP-AP-L Aberdeen Proving Ground, MD 21005-5066			10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARL-TR-695	
11. SUPPLEMENTARY NOTES *P. J. Stevens and F. J. Develin are employed by the University of Southern California. **M. J. Frisch is employed by Lorentzian Inc.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The unpolarized absorption and circular dichroism spectra of the fundamental vibrational transitions of the chiral molecule, 4-methyl-2-oxetanone, are calculated ab initio. Harmonic force fields are obtained using Density Functional Theory (DFT), MP2, and SCF methodologies and a [5s4p2d/3s2p] (TZ2P) basis set. DFT calculations use the Local Spin Density Approximation (LSDA), BLYP, and Becke3LYP (B3LYP) density functionals. Mid-IR spectra predicted using LSDA, BLYP, and B3LYP force fields are of significantly different quality, the B3LYP force field yielding spectra in clearly superior, and overall excellent, agreement with experiment. The MP2 force field yields spectra in slightly worse agreement with experiment than the B3LYP force field. The SCF force field yields spectra in poor agreement with experiment. The basis set dependence of B3LYP force fields is also explored: the 6-31G* and TZ2P basis sets give very similar results while the 3-21G basis set yields spectra in substantially worse agreement with experiment.				
14. SUBJECT TERMS vibration circular dichroism spectra, density functional theory, MP2 force fields			15. NUMBER OF PAGES 24	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

INTENTIONALLY LEFT BLANK.

ACKNOWLEDGMENTS

The past and ongoing support of studies of Vibrational Circular Dichroism at the University of Southern California (USC) by the National Science Foundation (NSF), the National Institutes of Health (NIH), the North Atlantic Treaty Organization (NATO), and the San Diego Supercomputer Center is gratefully acknowledged. Dr. C. F. Chabalowski would like to thank the U.S. Army Edgewood Research, Development, and Engineering Center (ERDEC) for partial support of this work.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution _____	
Availability Codes	
Dist	Avail and/or Special
A-1	

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGMENTS	iii
LIST OF FIGURES	vii
1. INTRODUCTION	1
2. METHODS	2
3. RESULTS AND DISCUSSION	4
4. CONCLUSION	9
5. REFERENCES	13
DISTRIBUTION LIST	15

INTENTIONALLY LEFT BLANK.

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1.	(a) Experimental unpolarized absorption spectrum of 1 (in CCl ₄ and CS ₂ solutions, from Devlin et al. [to be published]);	
	(b) Fundamental unpolarized absorption spectrum of 1 (obtained from (a) by Lorentzian fitting [Devlin et al., to be published]);	
	(c)–(g) Calculated unpolarized absorption spectrum of 1 (In (c), (d), and (e), force fields and APTs are calculated using DFT and B3LYP, BLYP, and LSDA density functionals, respectively. In (f) and (g), force fields and APTs are calculated using MP2 and SCF methodologies. All calculations use the TZ2P basis set. Lorentzian band half-widths, γ , are 4.0 cm ⁻¹ .)	5
2.	(a) Experimental VCD spectrum of R- 1 (in CCl ₄ and CS ₂ solutions, from Devlin et al. [to be published]);	
	(b) Fundamental VCD spectrum of R- 1 (obtained from (a) by Lorentzian fitting [Devlin et al., to be published]);	
	(c)–(g) Calculated VCD spectrum of R- 1 (These originate as in Figure 1. All calculations use SCF TZ2P local AATs. Lorentzian band half-widths, γ , are 4.0 cm ⁻¹ .)	6
3.	(a) Fundamental unpolarized absorption spectrum of 1 (as in Figure 1b);	
	(b)–(d) Calculated unpolarized absorption spectrum of 1 (In (b), (c), and (d), force fields and APTs are calculated using DFT, the B3LYP density functional and, respectively, the TZ2P, 6-31G* and 3-21G Basis sets. Lorentzian band half-widths, γ , are 4.0 cm ⁻¹ .)	10
4.	(a) Fundamental VCD spectrum of R- 1 (as in Figure 2b);	
	(b)–(d) Calculated VCD spectrum of R- 1 (These originate as in Figure 3. All calculations use SCF TZ2P local AATs. Lorentzian band half-widths, γ , are 4.0 cm ⁻¹ .)	11

INTENTIONALLY LEFT BLANK.

1. INTRODUCTION

The use of density functional theory (DFT) in the *ab initio* calculation of molecular properties has recently increased dramatically (Ziegler 1991; Labanowski and Andzelm [editors] 1991). This can be attributed to the following: (1) the development of new and more accurate density functionals; (2) the increasing versatility, efficiency, and availability of DFT codes; and, most importantly, (3) the superior ratio of accuracy to effort exhibited by DFT computations relative to other *ab initio* methodologies.

We focus here on the application of DFT to the prediction of vibrational spectra. With the most accurate density functionals currently in use, DFT predicts harmonic vibrational frequencies of substantially higher accuracy than obtained via SCF calculations and of similar accuracy to the predictions of MP2 calculations (Johnson, Gill, and Pople 1993; Handy, Murray, and Amos 1993). Since DFT calculations are substantially less demanding computationally than MP2 calculations, it is clear that DFT offers significant advantages in predicting harmonic vibrational force fields, frequencies, and spectra.

In particular, DFT promises to greatly facilitate the calculation of Vibrational Circular Dichroism (VCD) spectra (Stephens and Lowe 1985). Experience has shown that (with the exception of small, symmetrical molecules) the SCF harmonic force fields are generally of insufficient accuracy to provide useful predictions of VCD spectra (Kawiecki 1988; Jalkanen 1989; Bursi 1991). Calculations using MP2 harmonic force fields are substantially more successful (Bursi 1991; Amos, Handy, and Palmieri 1990; Stephens et al. 1993, Stephens et al., to be published; Devlin and Stephens 1994), but also much more demanding. The lower computational demands of DFT suggest that DFT may become preferred over the MP2 methodology in the future *ab initio* predictions of VCD spectra.

We report here calculations of the unpolarized absorption and circular dichroism spectra arising from the fundamental vibrational excitations of the chiral molecule 4-methyl-2-oxetanone, **1**, previously studied (both experimentally and theoretically) in our laboratories (Jalkanen 1989; Devlin et al., to be published). We compare the accuracies of spectra predicted using DFT, MP2, and SCF harmonic force fields. The accuracy of DFT calculations is dependent on the density functional employed. In this work, we compare three significantly different density functionals: Local Spin Density Approximation (LSDA), BLYP, and Becke3LYP. The LSDA and BLYP functionals have been widely used (Ziegler 1991; Labanowski and Andzelm [editors] 1991; Johnson, Gill, and Pople 1993; Handy, Murray, and Amos 1993). The Becke3LYP functional is a hybrid of several components, whose relative weights are chosen by reference

to experimental thermochemical data. The accuracies of such hybrid functionals in predicting molecular geometries and vibrational frequencies have not yet been thoroughly characterized.

Our calculations take advantage of recent developments in the computation of DFT and MP2 harmonic force fields via analytical derivative techniques (Trucks et al., unpublished; Johnson and Frisch 1993, 1994).

2. METHODS

Harmonic force fields for **1** were calculated ab initio at SCF and MP2 levels and using DFT via GAUSSIAN 92 and GAUSSIAN 92/DFT (Frisch et al. 1992, 1993). All force fields were calculated using analytical derivative methods (Trucks et al., unpublished; Johnson and Frisch 1993, 1994). DFT calculations were carried out using three density functionals:

- (1) LSDA (Local Spin Density Approximation). This uses the standard local exchange functional (Becke 1989) and the local correlation functional of Vosko, Wilk, and Nusair (VWN) (1980).
- (2) BLYP. This combines the standard local exchange functional with the gradient correction of Becke (1989) and uses the Lee-Yang-Parr (1988) correlation functional (which also includes density gradient terms).
- (3) Becke3LYP. This functional is a hybrid of exact (Hartree-Fock) exchange with local and gradient-corrected exchange and correlation terms, as first suggested by Becke (1993). The exchange-correlation functional proposed and tested by Becke was:

$$E_{xc} = (1 - a_0)E_x^{LSDA} + a_0E_x^{HF} + a_x\Delta E_x^{B88} + E_c^{LSDA} + a_c\Delta E_c^{PW91} \quad (1)$$

Here ΔE_x^{B88} is Becke's gradient correction to the exchange functional and ΔE_c^{PW91} is the Perdew-Wang gradient correction to the correlation functional (Perdew 1991). Becke suggested coefficients $a_0 = 0.2$, $a_x = 0.72$, and $a_c = 0.81$, based on fitting to heats of formation of small molecules. Only single-point energies were involved in the fit; no molecular geometries or frequencies were used. The Becke3LYP functional in Gaussian 92/DFT uses the values of a_0 , a_x , and a_c suggested by Becke, but uses LYP for the correlation functional. Since LYP does not have an easily separable local component, the VWN local correlation expression has been used to provide the different coefficients of local and gradient corrected correlation functionals:

$$E_{xc}^{B3LYP} = (1 - a_0)E_x^{LSDA} + a_0E_x^{HF} + a_x\Delta E_x^{B88} + a_cE_c^{LYP} + (1 - a_c)E_c^{VWN} . \quad (2)$$

The standard fine grid in Gaussian 92/DFT (Trucks and Frisch, to be published) was used in all DFT calculations. This grid was produced from a basic grid having 75 radial shells and 302 angular points per radial shell, leaving about 7,000 points per atom while retaining similar accuracy to the original (75,302) grid. Becke's (1988) numerical integration techniques were employed.

Atomic Polar Tensors (APTs) (Stephens et al. 1990) were calculated at the same time as harmonic force fields. Atomic Axial Tensors (AATs) were calculated using the Distributed Origin gauge (Stephens et al. 1990; Stephens 1987), in which the AAT of nucleus λ , $(M_{\alpha\beta}^{\lambda})^0$, with respect to origin O is given by

$$(M_{\alpha\beta}^{\lambda})^0 = (I_{\alpha\beta}^{\lambda})^{\lambda} + \frac{i}{4\hbar c} \sum_{\gamma\delta} \epsilon_{\beta\gamma\delta} R_{\lambda\gamma}^0 P_{\alpha\delta}^{\lambda} \quad (3)$$

where R_{λ}^0 is the equilibrium position of nucleus λ relative to origin O, $P_{\alpha\beta}^{\lambda}$ is the APT of nucleus λ , and $(I_{\alpha\beta}^{\lambda})^{\lambda}$ is the electronic AAT of nucleus λ calculated with the origin at R_{λ}^0 . "Local" AATs, $(I_{\alpha\beta}^{\lambda})^{\lambda}$, are calculated using CADPAC (version 5) (Amos 1993) at the SCF level. (At the present time, neither MP2 nor DFT codes for AATs are available).

Vibrational frequencies, dipole strengths, and rotational strengths were calculated from the harmonic force fields, APTs, and AATs (Stephens 1985; Jalkanen et al. 1990; Kawiecki et al. 1991). Unpolarized absorption and circular dichroism spectra were synthesized thence using Lorentzian band shapes (Kawiecki et al. 1988). Calculations used 3-21G, 6-31G* (Hehre et al. 1986) and [5s4p2d/3s2p], TZ2P (Stephens et al. 1990) basis sets (66, 102, and 228 basis functions, respectively). SCF and MP2 calculations were carried out using the San Diego Supercomputer Center CRAY-YMP and C90 machines. DFT calculations were carried out at Lorentzian Inc. using IBM RS/6000-590 and Silicon Graphics Incorporated (SGI) Challenge machines.

3. RESULTS AND DISCUSSION

Unpolarized absorption and circular dichroism spectra of **1**, measured in CS₂ and CCl₄ solutions (Devlin et al., to be published), are reproduced in Figures 1 and 2 over the spectral range 650–1,500 cm⁻¹. Frequencies, dipole strengths, and rotational strengths of transitions assigned to fundamental excitations, obtained by Lorentzian fitting of the experimental spectra (Devlin et al., to be published), are given in Table 1. The spectra attributable to these transitions alone are also shown in Figures 1 and 2.

Table 1. Frequencies, Dipole Strengths, and Rotational Strengths of **1**^a

MP2			DFT/B3LYP			EXPT ^b		
ν	D	R	ν	D	R	ν	D	R
1,528	14	-5	1,500	13	-5	1,453	25	-2
1,516	8	4	1,489	8	5	1,441	3	3
1,479	47	-11	1,462	43	-11	1,419	35	-9
1,438	52	11	1,422	51	11	1,387	58	13
1,394	52	27	1,389	50	28	1,350	40	42
1,320	97	9	1,316	112	3	1,284	161	26
1,230	47	-11	1,229	14	-4	1,198	47	-21
1,216	14	0	1,204	19	-5	1,178	35	-17
1,146	191	82	1,137	327	99	1,118	483	178
1,125	219	-40	1,113	87	-58	1,099	51	-83
1,077	10	-12	1,076	11	-6	1,055	40	-23
1,056	271	-50	1,041	268	-50	1,022	251	-102
973	126	10	966	91	7	959	70	12
914	9	-29	914	8	-27	896	10	-54
873	221	33	850	310	46	836	371	78
829	97	26	817	34	10	812	55	29
714	3	-3	712	2	2	711	13	15

^a Frequencies, ν , in cm⁻¹; dipole strengths D in 10⁻⁴⁰esu²cm²; rotational strengths R in 10⁻⁴⁴esu²cm².

R values are for R-(+)-**1**.

^b Devlin et al., to be published.




Figure 1. (a) Experimental unpolarized absorption spectrum of 1 (in CCl₄ and CS₂ solutions, from Devlin et al. [to be published]);

(b) Fundamental unpolarized absorption spectrum of 1 (obtained from (a) by Lorentzian fitting [Devlin et al., to be published]);

(c)–(g) Calculated unpolarized absorption spectrum of 1 (In (c), (d), and (e), force fields and APTs are calculated using DFT and B3LYP, BLYP, and LSDA density functionals, respectively. In (f) and (g), force fields and APTs are calculated using MP2 and SCF methodologies. All calculations use the TZ2P basis set. Lorentzian band half-widths, γ , are 4.0 cm⁻¹).




Figure 2. (a) Experimental VCD spectrum of R-1 (in CCl_4 and CS_2 solutions, from Devlin et al. [to be published]);
 (b) Fundamental VCD spectrum of R-1 (obtained from (a) by Lorentzian fitting [Devlin et al., to be published]);
 (c)–(g) Calculated VCD spectrum of R-1 (These originate as in Figure 1. All calculations use SCF TZ2P local AATs. Lorentzian band half-widths, γ , are 4.0 cm^{-1} .).

Harmonic frequencies, dipole strengths, and rotational strengths have been calculated for **1** using DFT at the TZ2P basis set level and utilizing the LSDA, BLYP, and B3LYP density functionals. Unpolarized absorption spectra and VCD spectra derived thence are displayed in Figures 1 and 2. The absorption and VCD spectra predicted using the LSDA, BLYP, and B3LYP functionals are substantially different, demonstrating that the choice of functional is of importance. Comparison to the fundamental absorption and VCD spectra shows that the B3LYP functional provides spectra in best agreement with experiment when judged on the basis of the qualitative pattern of frequencies and intensities. The BLYP and LSDA functionals yield spectra in significantly worse agreement. The variations among predicted spectra originate predominantly in the variations in the harmonic force fields (variations in the APTs are of minor consequence in comparison). We therefore conclude that the B3LYP, BLYP, and LSDA force fields are of significantly different accuracy, the B3LYP force field being the most accurate.

The qualitative agreement of the absorption and VCD spectra calculated using the B3LYP functional with the experimental spectra is impressive. The quantitative agreement of theory and experiment can be assessed from the calculated and experimental frequencies, dipole strengths, and rotational throughout, the average and maximum deviations being $25\text{ cm}^{-1}/2.0\%$ and $48\text{ cm}^{-1}/3.3\%$. Some fraction of the differences in calculated and experimental frequencies can, of course, be attributed to anharmonicity and solvent effects. The calculated dipole strengths are in excellent qualitative agreement with experimental values; quantitative differences are attributable to a combination of residual errors in the DFT/B3LYP/TZ2P calculation, anharmonicity and solvent effects, and to experimental error. Calculated rotational strengths are also in excellent qualitative agreement with experimental values. The quantitative agreement is somewhat poorer than in the case of dipole strengths. This is undoubtedly attributable to the lower accuracy of the AATs, compared to the APTs, due to the absence of correlation in the calculation of local AATs.

The results of TZ2P MP2 calculations (Chabalowski, Devlin, and Stephens to be submitted) are also given in Table 1 and in Figures 1 and 2. The agreement with experiment of the spectra calculated using the MP2 force field is slightly worse than obtained from the B3LYP force field. The most prominent difference is in the relative absorption intensities of the bands calculated between $1,100$ and $1,150\text{ cm}^{-1}$. On the other hand, the agreement is significantly better than obtained from the BLYP and LSDA force fields. We conclude that the MP2 force field is respectively lower, higher, and higher in accuracy than the B3LYP, BLYP, and LSDA force fields.

The quantitative agreement of frequencies, dipole strengths, and rotational strengths calculated from the MP2 force field with experimental values can be assessed from Table 1. Calculated frequencies are in all cases greater than experimental frequencies, the average and maximum deviations being $36\text{ cm}^{-1}/3.0\%$ and $75\text{ cm}^{-1}/5.2\%$. The differences are in all cases greater than for the B3LYP force field, and, consequently, so are the average and maximum deviations. Calculated dipole strengths are, in most cases, further from experimental values than those obtained from the B3LYP force field. Comparison of both frequencies and dipole strengths thus support the conclusion that the harmonic force field is less accurately calculated by the MP2 methodology. Rotational strengths calculated from the MP2 force field are in good qualitative agreement with both experimental values and the predictions of the B3LYP force field. Neither of the calculations exhibits clear-cut quantitative superiority.

Absorption and VCD spectra obtained from the SCF harmonic force field are also shown in Figures 1 and 2. Both absorption and VCD spectra are qualitatively different from the experimental spectra, the differences being sufficiently large that a one-to-one correspondence between calculated and observed bands cannot be arrived at; i.e., the predictions are insufficiently accurate to yield a convincing assignment of the experimental spectra. It is obvious that the SCF force field is drastically lower in accuracy than the DFT and MP2 force fields.

The calculations presented in Figures 1 and 2 and in Table 1 use a large basis set—TZ2P—in order to reduce basis set errors to the maximum extent. Having established the superiority of the B3LYP functional over the LSDA and BLYP functionals, it is of interest to examine the basis set dependence of the spectra predicted using the B3LYP functional. Figures 3 and 4 compare absorption and VCD spectra calculated using the 3-21G and 6-31G* basis sets to those obtained at the TZ2P level and to experiment. Local AATs are calculated at the TZ2P basis set level throughout, and the variations in spectra are due entirely to variations in the DFT force fields and APTs with basis sets. The absorption and VCD spectra predicted using the 6-31G* and TZ2P basis sets are identical qualitatively and very little different quantitatively. The agreement with the experimental spectra is almost equally good. The 3-21G basis set yields spectra substantially different from those obtained using the 6-31G* and TZ2P basis sets and in much worse agreement with experiment.

4. CONCLUSION

In the case of **1**, and the TZ2P basis set level, the DFT force field calculated using the B3LYP density functional yields mid-infrared (IR) absorption and VCD spectra in impressive agreement with experiment. DFT force fields calculated using the LSDA and BLYP density functionals yield spectra of significantly lower accuracy. The MP2 and SCF force fields yield spectra slightly less and much less accurate, respectively, than the DFT/B3LYP force field. Calculations on a wide variety of molecules are clearly required to define the generality of these results. Such calculations are under way.

While the LSDA and BLYP functionals have been widely used in DFT calculations of vibrational frequencies (Johnson, Gill, and Pople 1993; Handy, Murray, and Amos 1993), this is not the case for the hybrid functionals recently introduced by Becke (1993). Since these functionals incorporate parameters determined by fitting to experimental thermochemical data, their accuracy in predicting molecular geometries and vibrational force fields was not obvious. Our results for **1** using the B3LYP functional support the conclusion that the methodology of Becke leads to molecular force fields, as well as thermochemical properties, of improved accuracy. Our results also show that B3LYP calculations converge rapidly with increasing basis set size and that the cost-to-benefit ratio is optimal at the 6-31G* basis set level. 6-31G* will be the basis set of choice in B3LYP calculations on much larger molecules.

Our calculations clearly demonstrate the sensitivity of predictions of vibrational spectra to the accuracy of the force field, and, in the case of DFT calculations, to the choice of density functional. The comparison to experimental spectra of vibrational spectra predicted from future generations of density functionals should be of significant utility in defining their relative accuracies. In the case of chiral molecules, the information content of vibrational spectroscopy is considerably enhanced when absorption and VCD spectra are utilized in combination.

VCD spectroscopy is potentially a powerful tool in elucidating the stereochemistry of chiral molecules. The work reported here clearly supports the expectation that DFT now provides a basis for calculations of VCD spectra at a useful level of accuracy for molecules much larger than **1**. This capability will be documented in forthcoming publications.




Figure 3. (a) Fundamental unpolarized absorption spectrum of 1 (as in Figure 1b);

(b)–(d) Calculated unpolarized absorption spectrum of 1 (In (b), (c), and (d), force fields and APTs are calculated using DFT, the B3LYP density functional, and, respectively, the TZ2P, 6-31G*, and 3-21G Basis sets. Lorentzian band half-widths, γ , are 4.0 cm^{-1}).




Figure 4. (a) Fundamental VCD spectrum of R-1 (as in Figure 2b);

(b)–(d) Calculated VCD spectrum of R-1 (These originate as in Figure 3. All calculations use SCF TZ2P local AATs. Lorentzian band half-widths, γ , are 4.0 cm^{-1}).

INTENTIONALLY LEFT BLANK.

5. REFERENCES

- Amos, R. D., and J. E. Rice. "CADPAC: The Cambridge Analytic Derivatives Package," issue 5. Cambridge University, Cambridge, England, 1992.
- Amos, R., N. C. Handy, and P. Palmieri. Journal of Chemical Physics, vol. 93, p. 5796, 1990.
- Becke, A. D. Journal of Chemical Physics, vol. 88, p. 2547, 1988.
- Becke, A. D. in The Challenge of d and f Electrons: Theory and Computation, ch. 12, pp. 165-179. Edited by D. R. Salahub and M. C. Zerner. American Chemical Society, Washington, DC, 1989.
- Becke, A. D. Journal of Chemical Physics, vol. 98, p. 1372, 1993.
- Bursi, R. Ph.D. Thesis, University of Southern California, 1991.
- Chabalowski, C. F., F. J. Devlin, and P. J. Stephens. "Ab Initio Calculations of Vibrational Circular Dichroism and Infrared Spectra Using MP2 Force Fields: 4-Methyl-2-Oxetanone." U.S. Army Research Laboratory, Aberdeen Proving Ground, MD, to be published.
- Devlin, F., K. J. Jalkanen, T. Polonski, and P. J. Stephens, to be published.
- Devlin, F. J., and P. J. Stephens. Journal of the American Chemical Society, vol. 116, p. 5003, 1994.
- Frisch, M. J., G. W. Trucks, M. Head-Gordon, F. M. W. Gill, M. W. Wong, J. B. Foresman, B. G. Johnson, H. B. Schlegel, M. A. Robb, E. S. Replogle, R. Gomperts, J. L. Andres, K. Raghavachari, J. S. Binkley, C. Gonzalez, R. L. Martin, D. J. Fox, D. J. Defrees, J. Baker, J. J. P. Stewart, and J. A. Pople. "GAUSSIAN 92," Revision G. Gaussian, Inc., Pittsburg, PA, 1992.
- Frisch, M. J., G. W. Trucks, M. Head-Gordon, F. M. W. Gill, M. W. Wong, J. B. Foresman, B. G. Johnson, H. B. Schlegel, M. A. Robb, E. S. Replogle, R. Gomperts, J. L. Andres, K. Raghavachari, J. S. Binkley, C. Gonzalez, R. L. Martin, D. J. Fox, D. J. Defrees, J. Baker, J. J. P. Stewart, and J. A. Pople. "GAUSSIAN 92/DFT." Gaussian, Inc., Pittsburg, PA, 1993.
- Handy, N. C., C. W. Murray, and R. D. Amos. Journal of Physical Chemistry, vol. 97, p. 4392, 1993.
- Hehre, W., L. Random, P. R. Schleyer, and J. A. Pople. Ab Initio Molecular Orbital Theory. New York: Wiley, 1986.
- Jalkanen, K. J. Ph.D. Thesis, University of Southern California, 1989.
- Jalkanen, K. J., R. W. Kawiecki, P. J. Stephens, and R. D. Amos. Journal of Physical Chemistry, vol. 94, p. 7040, 1990.
- Johnson, B. G., and M. J. Frisch. Chemical Physics Letters, vol. 216, p. 133, 1993.
- Johnson, B. G., and M. J. Frisch. Journal of Chemical Physics, vol. 100, p. 7429, 1994.
- Johnson, B. G., P. M. W. Gill, and J. A. Pople. Journal of Chemical Physics, vol. 98, p. 5612, 1993.

- Kawiecki, R. W. Ph.D. Thesis, University of Southern California, 1988.
- Kawiecki, R. W., F. J. Devlin, P. J. Stephens, R. D. Amos, and N. C. Handy. Chemical Physics Letters, vol. 145, p. 411, 1988.
- Kawiecki, R. W., F. J. Devlin, P. J. Stephens, and R. D. Amos. Journal of Physical Chemistry, vol. 95, p. 9817, 1991.
- Labanowski, J. K., and J. Andzelm (editors). Density Functional Methods in Chemistry. New York, Springer-Verlag, 1991.
- Lee, C., W. Yang, and R. G. Parr. Physical Review B, vol. 37, p. 785, 1988.
- Perdew, J. P. in Electronic Structure of Solids. Edited by P. Ziesche and H. Eschrig, Berlin: Akademie Verlag, 1991.
- Stephens, P. J. Journal of Physical Chemistry, vol. 89, p. 748, 1985.
- Stephens, P. J. Journal of Physical Chemistry, vol. 91, p. 1712, 1987.
- Stephens, P. J., C. F. Chabalowski, F. J. Devlin, and K. J. Jalkanen. Chemical Physics Letters, vol. 225, p. 247, 1994.
- Stephens, P. J., K. J. Jalkanen, R. D. Amos, P. Lazzeretti, and R. Z. Zanasi. Journal of Physical Chemistry, vol. 94, p. 1811, 1990.
- Stephens, P. J., K. J. Jalkanen, F. J. Devlin, and C. F. Chabalowski. Journal of Physical Chemistry, vol. 97, p. 6107, 1993.
- Stephens, P.J., and M. A. Lowe. Annual Review of Physical Chemistry, vol. 36, p. 213, 1985.
- Trucks, G. W., and M. J. Frisch, to be published.
- Trucks, G. W., M. J. Frisch, M. Head-Gordon, J. Andres, J. B. Schlegel, and A. Salter, to be published.
- Vosko, S. H., L. Wilk, and M. Nusair. Canadian Journal of Physics, vol. 58, p. 1200, 1980.
- Ziegler, T. Chemical Reviews vol. 91, p. 65, 1991.

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>	<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
2	ADMINISTRATOR ATTN DTIC DDA DEFENSE TECHNICAL INFO CTR CAMERON STATION ALEXANDRIA VA 22304-6145	1	COMMANDER ATTN AMSMI RD CS R DOC US ARMY MISSILE COMMAND REDSTONE ARSNL AL 35898-5010
1	COMMANDER ATTN AMCAM US ARMY MATERIEL COMMAND 5001 EISENHOWER AVE ALEXANDRIA VA 22333-0001	1	COMMANDER ATTN AMSTA JSK ARMOR ENG BR US ARMY TANK AUTOMOTIVE CMD WARREN MI 48397-5000
1	DIRECTOR ATTN AMSRL OP SD TA US ARMY RESEARCH LAB 2800 POWDER MILL RD ADELPHI MD 20783-1145	1	DIRECTOR ATTN ATRC WSR USA TRADOC ANALYSIS CMD WSMR NM 88002-5502
3	DIRECTOR ATTN AMSRL OP SD TL US ARMY RESEARCH LAB 2800 POWDER MILL RD ADELPHI MD 20783-1145	1	COMMANDANT ATTN ATSH CD SECURITY MGR US ARMY INFANTRY SCHOOL FT BENNING GA 31905-5660
1	DIRECTOR ATTN AMSRL OP SD TP US ARMY RESEARCH LAB 2800 POWDER MILL RD ADELPHI MD 20783-1145		<u>ABERDEEN PROVING GROUND</u>
2	COMMANDER ATTN SMCAR TDC US ARMY ARDEC PCTNY ARSNL NJ 07806-5000	2	DIR USAMSAA ATTN AMXSY D AMXSY MP H COHEN
1	DIRECTOR ATTN SMCAR CCB TL BENET LABORATORIES ARSENAL STREET WATERVLIET NY 12189-4050	1	CDR USATECOM ATTN AMSTE TC
1	DIR USA ADVANCED SYSTEMS ATTN AMSAT R NR MS 219 1 R&A OFC AMES RESEARCH CENTER MOFFETT FLD CA 94035-1000	1	DIR USAERDEC ATTN SCBRD RT
		1	CDR USACBDCOM ATTN AMSCB CII
		1	DIR USARL ATTN AMSRL SL I
		5	DIR USARL ATTN AMSRL OP AP L

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	HQDA ATTN SARD TT DR F MILTON WASH DC 20310-0103
1	HQDA ATTN SARD TT MR J APPEL WASH DC 20310-0103
1	HQDA ATTN SARD TR MS K KOMINOS WASH DC 20310 0103
1	HQDA ATTN SARD TR DR R CHAIT WASH DC 20310 0103
1	HQDA ATTN OASA RDA DR C H CHURCH PENTAGON RM 3E486 WASH DC 20310-0103
4	COMMANDER ATTN R GHIRARDELLI D MANN R SINGLETON R SHAW US ARMY RSRCH OFFICE PO BOX 12211 RSRCH TRIGL PK NC 27709-2211
1	DIRECTOR ATTN AMXRO MCS K CLARK US ARMY RSRCH OFFICE PO BOX 12211 RSRCH TRIGL PK NC 27709-2211
1	DIRECTOR ATTN AMXRO RT IP LIBRARY SERVICES US ARMY RSRCH OFFICE PO BOX 12211 RSRCH TRIGL PK NC 27709-2211
2	COMMANDER ATTN SMCAR AEE B D S DOWNS US ARMY ARDEC PCTNY ARSNL NJ 07806-5000
2	COMMANDER ATTN SMCAR AEE J A LANNON US ARMY ARDEC PCTNY ARSNL NJ 07806-5000

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	COMMANDER ATTN SMCAR AEE BR L HARRIS US ARMY ARDEC PCTNY ARSNL NJ 07806-5000
2	COMMANDER ATTN AMSMI RD PR E A R MAYKUT AMSMI RD PR P R BETTS US ARMY MISSILE COMMAND REDSTONE ARSNL AL
1	OFFICE OF NAVAL RSCH DEPT OF THE NAVY ATTN R S MILLER CODE 432 800 N QUINCY ST ARLINGTON VA 22217
1	COMMANDER ATTN J RAMNARACE AIR 54111C NAVAL AIR SYSTEMS COMMAND WASH DC 20360
2	COMMANDER ATTN R BERNECKER R 13 G B WILMOT R 16 NAVAL SURFACE WARFARE CTR SLVR SPRNG MD 20903-5000
5	COMMANDER ATTN M C LIN J MCDONALD E ORAN J SHNUR R J DOYLE CODE 6110 NAVAL RSCH LAB WASH DC 20375
2	COMMANDER ATTN T BOGGS CODE 388 T PARR CODE 3895 NAVAL WEAPONS CTR CHINA LAKE CA 93555-6001
1	SUPERINTENDENT ATTN D W NETZER NAVAL POSTGRADUATE SCHOOL DEPT OF AERONAUTICS MONTEREY CA 93940

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
3	AL LSCF ATTN R CORLEY R GEISLER J LEVINE EDWARDS AFB CA 93523-5000
1	AFOSR ATTN J M TISHKOFF BOLLING AIR FORCE BASE WASH DC 20332
1	OSD SDIO IST ATTN L CAVENY PENTAGON WASH DC 20301-7100
1	COMMANDANT ATTN ATSF TSM CN USAFAS FT SILL OK 73503-5600
1	UNIV OF DAYTON RSRCH INSTITUTE ATTN D CAMPBELL AL PAP EDWARDS AFB CA 93523
1	NASA ATTN G B NORTHAM MS 168 LANGLEY RSCH CTR LANGLEY STATION HAMPTON VA 23365
4	NATIONAL BUREAU OF STANDARDS ATTN J HASTIE M JACOX T KASHIWAGI H SEMERJIAN US DEPT OF COMMERCE WASH DC 20234
2	DIRECTOR ATTN C WESTBROOK W TAO MS L 282 LAWRENCE LIVERMORE NATL LAB PO BOX 808 LIVERMORE CA 94550
1	DIRECTOR ATTN B NICHOLS T7 MS B284 LOS ALAMOS NATL LAB PO BOX 1663 LOS ALAMOS NM 87545

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	NATL SCIENCE FOUNDATION ATTN A B HARVEY WASH DC 20550
2	PRINCETON COMBUSTION RSRCH LAB INC ATTN N A MESSINA M SUMMERFIELD PRINCETON CORPORATE PLAZA BLDG IV STE 119 11 DEERPARK DR MONMOUTH JUNCTION NJ 08852
3	DIRECTOR ATTN S JOHNSTON P MATTERN D STEPHENSON SANDIA NATL LAB DIV 8354 LIVERMORE CA 94550
1	BRIGHAM YOUNG UNIV ATTN M W BECKSTEAD DEPT OF CHEMICAL ENGRG PROVO UT 84058
1	CA INSTITUTE OF TECH JET PROPULSION LAB ATTN L STRAND MS 125 224 4800 OAK GROVE DR PASADENA CA 91109
1	CALIFORNIA INSTITUTE OF TECHNOLOGY ATTN F E C CULICK MC 301 46 204 KARMAN LAB PASADENA CA 91125
1	UNIV OF CALIFORNIA ATTN LOS ALAMOS SCIENTIFIC LAB PO BOX 1663 MAIL STOP B216 LOS ALAMOS NM 87545
1	UNIV OF CALIFORNIA BERKELEY ATTN C BRADLEY MOORE CHEMISTRY DEPT 211 LEWIS HALL BERKELEY CA 94720
1	UNIV OF CALIFORNIA SAN DIEGO ATTN F A WILLIAMS AMES B010 LA JOLLA CA 92093

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
2	UNIV OF CALIFORNIA SANTA BARBARA ATTN K SCHOFIELD M STEINBERG QUANTUM INSTITUTE SANTA BARBARA CA 93106
1	UNIV OF COLORADO AT BOULDER ATTN J DAILY ENGRG CTR CAMPUS BOX 427 BOULDER CO 80309 0427
3	UNIV OF SOUTHERN CALIFORNIA ATTN R BEAUDET S BENSON C WITTIG DEPT OF CHEMISTRY LOS ANGELES CA 90007
1	CORNELL UNIV ATTN T A COOL DEPT OF CHEMISTRY BAKER LAB ITHACA NY 14853
1	UNIV OF DELAWARE ATTN T BRILL CHEMISTRY DEPT NEWARK DE 19711
1	UNIV OF FLORIDA ATTN J WINEFORDNER DEPT OF CHEMISTRY GAINESVILLE FL 32611
3	GEORGIA INSTITUTE OF TECHNOLOGY ATTN E PRICE W C STRAHLE B T ZINN SCHOOL OF AEROSPACE ENGRG ATLANTA GA 30332
1	UNIV OF ILLINOIS ATTN H KRIER DEPT OF MECH ENGRG 144MEB 1206 W GREEN ST URBANA IL 61801

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	THE JOHNS HOPKINS UNIV ATTN T W CHRISTIAN CHEMICAL PROPULSION INFORMATION AGENCY 10630 LITTLE PATUXENT PKWY STE 202 COLUMBIA MD 21044 3200
1	UNIV OF MICHIGAN ATTN G M FAETH GAS DYNAMICS LAB AEROSPACE ENGRG BLDG ANN ARBOR MI 48109 2140
1	UNIV OF MINNESOTA ATTN E FLETCHER DEPT OF MECHANICAL ENGRG MINNEAPOLIS MN 55455
4	PENNSYLVANIA STATE UNIV ATTN K KUO M MICCI S THYNELL V YANG DEPT OF MECHANICAL ENGRG UNIV PK PA 16802
1	POLYTECHNIC INSTITUTE OF NY ATTN S LEDERMAN GRADUATE CTR RTE 110 FARMINGDALE NY 11735
2	PRINCETON UNIV ATTN K BREZINSKY I GLASSMAN FORRESTAL CAMPUS LIB PO BOX 710 PRINCETON NJ 08540
1	PURDUE UNIV ATTN J R OSBORN SCHOOL OF AERONAUTICS & ASTRONAUTICS GRISSOM HALL WEST LAFAYETTE IN 47906
1	PURDUE UNIV ATTN E GRANT DEPT OF CHEMISTRY WEST LAFAYETTE IN 47906

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
2	PURDUE UNIV ATTN N M LAURENDEAU S N B MURTHY SCHOOL OF MECHANICAL ENGRG TSPC CHAFFEE HALL WEST LAFAYETTE IN 47906
1	RENSSELAER POLYTECHNIC INST ATTN A FONTIJN DEPT OF CHEMICAL ENGRG TROY NY 12181
1	STANFORD UNIV ATTN R HANSON DEPT OF MECHANICAL ENGRG STANFORD CA 94305
1	UNIV OF TEXAS ATTN W GARDINER DEPT OF CHEMISTRY AUSTIN TX 78712
1	VA POLYTECHNIC INST & STATE UNIV ATTN J A SCHETZ BLACKSBURG VA 24061
1	APPLIED COMBUSTION TECH INC ATTN A M VARNEY PO BOX 607885 ORLANDO FL 32860
2	APPLIED MECHANICS REVIEWS ATTN R E WHITE A B WENZEL THE AMERICAN SOCIETY OF MECH ENGRG 345 E 47TH ST NEW YORK NY 10017
1	ATLANTIC RSCH CORP ATTN R H W WAESCHE 7511 WELLINGTON RD GAINESVILLE VA 22065
1	TEXTRON DEFENSE SYSTEMS ATTN A PATRICK 2385 REVERE BEACH PKWY EVERETT MA 02149-5900

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	BATTELLE TWSTIAC HUGGINS 505 KING AVE COLUMBUS, OH 43201-2693
1	COHEN PROFESSIONAL SERVICES ATTN N S COHEN 141 CHANNING ST REDLANDS CA 92373
1	EXXON RSCH & ENGRG CO ATTN A DEAN RTE 22E ANNANDALE NJ 08801
1	GENERAL APPLIED SCIENCE LAB INC 77 RAYNOR AVE RONKONKAMA NY 11779-6649
1	GENERAL ELECTRIC ORD SYSTEMS ATTN J MANDZY 100 PLASTICS AVE PITTSFIELD MA 01203
1	GENERAL MOTORS RSCH LAB ATTN T SLOANE PHYSICAL CHEMISTRY DEPT WARREN MI 48090-9055
2	HERCULES INC ATTN W B WALKUP E A YOUNT ALLEGHENY BALLISTICS LAB ROCKET CTR WV 26726
1	HERCULES INC ATTN R V CARTWRIGHT 100 HOWARD BLVD KENVIL NJ 07847
1	ALLIANT TECHSYSTEMS INC ATTN D E BRODEN MS MN50 2000 MARINE SYSTEMS GROUP 600 2ND ST NE HOPKINS MN 55343

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	ALLIANT TECHSYSTEMS INC ATTN R E TOMPKINS MN 11 2720 600 SECOND ST N HOPKINS MN 55343
1	IBM CORPORATION ATTN A C TAM RSRCH DIV 5600 COTTLE RD SAN JOSE CA 95193
1	IIT RSRCH INSTITUTE ATTN R F REMALY 10 W 35TH ST CHICAGO IL 60616
1	LOCKHEED MISSILES & SPACE CO ATTN GEORGE LO DEPT 52 35 B204 2 3251 HANOVER ST PALO ALTO CA 94304
1	OLIN ORDNANCE ATTN V MCDONALD LIB PO BOX 222 ST MARKS FL 32355-0222
1	PAUL GOUGH ASSOCIATES INC ATTN P S GOUGH 1048 S ST PORTSMOUTH NH 03801-5423
1	HUGHES AIRCRAFT CO ATTN T E WARD 8433 FALLBROOK AVE CANOGA PK CA 91303
1	ROCKWELL INTERNATIONAL CORP ATTN J E FLANAGAN HB02 ROCKETDYNE DIV 6633 CANOGA AVE CANOGA PK CA 91304
1	SCIENCE APPLICATIONS INC ATTN R B EDELMAN 23146 CUMORAH CREST WOODLAND HILLS CA 91364

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
3	SRI INTERNATIONAL ATTN G SMITH D CROSLEY D GOLDEN 333 RAVENSWOOD AVE MENLO PK CA 94025
1	STEVENS INSTITUTE OF TECH ATTN R MCALEVY III DAVIDSON LAB HOBOKEN NJ 07030
1	SVERDRUP TECH INC ATTN R J LOCKE MS SVR 2 LERC GROUP 2001 AEROSPACE PKWY BROOK PK OH 44142
1	SVERDRUP TECH INC ATTN J DEUR 2001 AEROSPACE PKWY BROOK PK OH 44142
3	THIOKOL CORPORATION ATTN R BIDDLE R WILLER TECH LIB ELKTON DIV PO BOX 241 ELKTON MD 21921
3	THIOKOL CORP ATTN S J BENNETT WASATCH DIV PO BOX 524 BRIGHAM CITY UT 84302
1	UNITED TECH RSCH CTR ATTN A C ECKBRETH E HARTFORD CT 06108
1	UNITED TECH CORP ATTN R R MILLER CHEMICAL SYSTEM DIV PO BOX 49028 SAN JOSE CA 95161-9028
1	UNIVERSAL PROPULSION CO ATTN H J MCSPADDEN 25401 N CENTRAL AVE PHOENIX AZ 85027-7837

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	VERITAY TECH INC ATTN E B FISHER 4845 MILLERSPORT HWY E AMHERST NY 14051-0305
1	FREEDMAN ASSOCIATES ATTN E FREEDMAN 2411 DIANA RD BALTIMORE MD 21209-1525
6	ALLIANT TECHSYSTEMS ATTN J BODE C CANDLAND L OSGOOD R BURETTA R BECKER M SWENSON 600 SECOND ST NE HOPKINS MN 55343
1	US ARMY BENET LAB ATTN SAM SOPOK SMCAR CCB B WATERVLIET NY 12189

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
	<u>ABERDEEN PROVING GROUND</u>
36	DIR USARL ATTN AMSRL WT P, A W HORST AMSRL-WT-PC, R A FIFER G F ADAMS W R ANDERSON R A BEYER S W BUNTE C F CHABALOWSKI K P MCNEILL-BOONSTOPPEL A COHEN R CUMPTON R DANIEL D DEVYNCK N F FELL B E FORCH J M HEIMERL A J KOTLAR M R MANAA W F MCBRATNEY K L MCNESBY S V MEDLIN M S MILLER A W MIZIOLEK S H MODIANO J B MORRIS J E NEWBERRY S A NEWTON R PATEL R A PESCE RODRIGUEZ B M RICE R C SAUSA M A SCHROEDER J A VANDERHOFF M WENSING A WHREN J M WIDDER C WILLIAMSON

INTENTIONALLY LEFT BLANK.

USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. ARL Report Number ARL-TR-695 Date of Report February 1995

2. Date Report Received _____

3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) _____

4. Specifically, how is the report being used? (Information source, design data, procedure, source of ideas, etc.) _____

5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided, or efficiencies achieved, etc? If so, please elaborate. _____

6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) _____

CURRENT
ADDRESS

Organization

Name

Street or P.O. Box No.

City, State, Zip Code

7. If indicating a Change of Address or Address Correction, please provide the Current or Correct address above and the Old or Incorrect address below.

OLD
ADDRESS

Organization

Name

Street or P.O. Box No.

City, State, Zip Code

(Remove this sheet, fold as indicated, tape closed, and mail.)
(DO NOT STAPLE)